



EFFICACY OF *DELONIX REGIA* RAFIN (SYN. *POINCIANA REGIA* BOJER EX. HOOK) FOR POTENTIAL ANTIFUNGAL ACTIVITY

R. A. SHARMA*, P. CHANDRAWAT, S. SHARMA, D. SHARMA, B. SHARMA AND D. SINGH

Genetics and Biochemistry Laboratory, Department of Botany,

University of Rajasthan, Jaipur - 302 004

E-mail: sharma_ra2007@yahoo.co.in

KEY WORDS

Delonix regia
Aspergillus niger
A. flavus
R. bataticola
F. oxysporum

Received on :

05.05.2010

Accepted on :

21.07.2010

*Corresponding
author

ABSTRACT

Dry powdered materials of flowers and seeds of *D. regia* were sequentially solvent extracted with various solvents to isolate metabolite-rich fractions such as flavonoids, anthraquinones and sterols and also evaluate for antifungal activity against *Aspergillus niger*, *A. flavus*, *Rhizopus bataticola* and *Fusarium oxysporum* by agar disk diffusion method. In both plant parts, flavonoids showed maximum antifungal activity against *A. flavus*.

INTRODUCTION

Now a days multiple drug resistance has developed due to the indiscriminate use of commercial antimicrobial drugs commonly use in the treatment of infectious diseases (Davis, 1994; Service, 1995). But sometimes antibiotics are associated with adverse effects on the host including hypersensitivity, immune suppression and allergic reactions (Ahmad *et al.*, 1998). This situation forced scientists to search for new antimicrobial substances. Given the alarming incidence of antibiotic resistance in bacteria of medical importance (Monroe and Polk, 2000), there is a constant need for new and effective therapeutic agents (Bhavanani and Ballow, 2000). Therefore, there is a need to develop alternative antimicrobial drugs for the treatment of infectious diseases from medicinal plants (Clark, 1996; Cordell, 2000). There are several reports on the antimicrobial activity of different herbal extracts in different regions of the world (Chung *et al.*, 2004; Nair and Chandra, 2004; De Boer *et al.*, 2005; Nair *et al.*, 2005).

Plant based antimicrobials represent a vast untapped source of medicines and further exploration of plant antimicrobials need to occur. Antimicrobials of plant origin have enormous therapeutic potential. They are effective in the treatment of infectious diseases while simultaneously mitigating many of the side-effects that are often associated with synthetic antimicrobials (Wu *et al.*, 1999).

In *D. regia*, these compounds are mostly secondary metabolites such as flavonoids (Subramanian *et al.*, 1966) sterols (Guerere *et al.*, 1986) tannins and phenol compounds (E. Sherbeing *et al.*, 1971). Some carotenoids like β -carotene, zeaxanthin etc.

(Jungalwala and Cama, 1962; Sun Gene and Co., 2004; Krishna and Grampurohit, 2005) also exhibits antimicrobial activities. In the present work *Delonix regia* Rafin (syn. *Poinciana regia* Bojer ex. Hook) was evaluated for its antifungal properties against some pathogenic fungi.

MATERIALS AND METHODS

Plant collection

The plant was collected from campus of University of Rajasthan, Jaipur, India, and authenticated from the herbarium, Department of Botany, University of Rajasthan, Jaipur, India. Fresh plant material was washed under running tap water, air dried and then homogenized to fine powder and stored in airtight bottles.

Source of test organisms

The pure cultures of test fungi, namely *Aspergillus niger*, *A. flavus*, *Rhizopus bataticola* and *Fusarium oxysporum* were obtained from the Seed Pathology Laboratory, Department of Botany, University of Rajasthan, Jaipur, India, which were maintained on Potato Dextrose Agar medium.

Culture of test microbes

For the cultivation of fungi, Potato Dextrose Agar Medium (PDA) was prepared by mixing 1000 ml. v. potato infusion + 20g agar + 20g glucose, followed by autoclaving and then test fungi were incubated at $27^{\circ}\text{C} \pm 1^{\circ}\text{C}$ for 48h and the cultures were maintained on the same medium by regular subculturing. To prepare the test plates, 10-15 ml. of the respective medium was poured into the Petri dishes and used for screening. For assessing the fungicidal efficacy, a uniform spread of the test

Table 1: Fungicidal efficacy of extracts of *Debinix regia*

Fungi organism	Extracts of dried flower				Extracts of dried seed material													
	PE	C ₆ H ₆	CH ₂ COH	CHCl ₃	EROH	AQ	FLAV.	STER.	ANTHR.	PE	C ₆ H ₆	CH ₂ COH	CHCl ₃	EROH	AQ	FLA.	STE.	ANT.
<i>A. flavus</i>	IZ	18.00	16.00	23.00	21.00	±	23.00	24.00	24.00	16.00	14.00	11.00	14.00	±	-	22.00	13.00	16.00
	AI*	0.90	0.80	1.15	1.05	1.05	1.25	1.20	1.20	0.80	0.70	0.55	0.70	±	-	1.10	0.65	0.80
<i>A. niger</i>	IZ	15.00	18.00	22.00	10.00	18.00	22.00	20.00	18.00	14.00	15.00	±	16.00	-	-	16.00	21.00	19.00
	AI	0.76	0.85	1.04	0.47	0.85	1.04	0.95	0.85	0.66	0.71	±	0.76	-	-	0.76	1.00	0.90
<i>R. bataticola</i>	IZ	18.00	17.00	18.00	-	-	23.00	24.00	19.00	15.00	16.00	18.00	±	14.00	13.00	21.00	21.00	19.00
	AI	0.72	0.56	0.72	-	-	0.92	0.96	0.76	0.60	0.64	0.72	±	0.56	0.52	0.84	0.84	0.76
<i>F. moniliforme</i>	IZ	9.00	9.00	11.00	±	±	14.00	13.00	13.00	±	13.00	7.00	8.00	8.00	19.00	17.00	14.00	
	AI	1.52	1.12	1.37	±	±	1.75	1.62	1.62	1.62	1.62	0.87	1.00	1.00	2.37	2.12	1.75	

Iz: Inhibitory zone (mm) including the diameter of disc; AI: Inhibitory index; H: Not measurable activity

*Activity index (AI) =

Inhibition area of the wet sample
Inhibition area of the standard

Abbreviations: PE = Petroleum ether; C₆H₆ = Benzene; CH₂COH = Acetone; CHCl₃ = Chloroform; EROH = Ethanol; AQ = Aqueous; FLAV. = Flavonoid; STER. = Sterol; ANTHR. = Anthracene; PE = Petroleum ether; P.A. = Flavonoid; STE = Sterol; Standard: Chitinazolin = 100 (µg/ml) disc




Plate 1: A. *Aspergillus niger*; B. *Aspergillus flavus*; C. *Rhizopus bataticola*; D. *Fusarium moniliforme*
a. Petroleum ether; b. Benzene; c. Acetone; d. Chloroform; e. Alcohol and f. Water.




Plate 2: E. *Aspergillus niger*; F. *Aspergillus flavus*; G. *Rhizopus bataticola*; H. *Fusarium moniliforme*
a. Petroleum ether; b. Benzene; c. Acetone; d. Chloroform; e. Alcohol and f. Water.




Plate 3: I. *Aspergillus niger*; J. *Aspergillus flavus*; K. *Rhizopus bataticola*; L. *Fusarium moniliforme*

a. Petroleum ether; b. Benzene; c. Acetone; d. Chloroform; e. Alcohol and f. Water.

fungi was made using the sterile swab.

Preparation of test extract

Dry and powdered materials of flowers and seeds (50 g each) of *D. regia* were sequentially soxhlet extracted with petroleum ether (600-800C), benzene, acetone, chloroform, alcohol and water successively, similarly, various metabolites-rich fractions, such as flavonoids, anthraquinones and sterols were also extracted separately by using established protocols.

All these extracts/fractions were stored at 90C in refrigerator until screened, when their final volume was raised to a known concentration (1g/mL) in their respective solvents before use.

Antifungal assay

The antimicrobial activity of different plant species was evaluated by agar disk diffusion method (Gould and Bowie, 1952; Bauer et al., 1966; Sallie et al., 1996) The different test organisms were preceded separately using a sterile swab over previously sterilized culture medium plates and the zones of inhibition were measured around sterilized dried discs of Whatman No. 1 paper (6 mm in diameter), which were containing 4 mg per disc (or 0.4 mL) of the test extracts of control (0.4 g/mL of the respective solvent) or Clotrimazole (100 units/mL) as reference drugs separately. Such treated discs were air dried at room temperature to remove any residual solvent which might interfere with the determination. These plates were initially placed at low temperature for 1 h, so as to allow the maximum diffusion of the compounds from the test discs into the agar plates and later, incubated at 370C for 48 h, after which the zones of inhibition could be easily observed. Five replicates of each test extract were examined and the mean values were then referred.

RESULTS AND DISCUSSION

The result of antifungal activity of *D. regia*, are cited in the Table 1. The extracts of sequential extraction with various solvents namely - pt. ether, benzene, acetone, chloroform, ethanol, aqueous and flavonoids, anthraquinones and sterols were used against certain pathogenic fungi namely *F. moniliforme*, *Rhizopus bataticola*, *Aspergillus niger* and *A. flavus*. Although all the bioactive compounds - the secondary metabolites flavonoids, anthraquinones and sterols exhibits antifungal activity, but flavonoids extracted from flowers (IZ : 23mm; AI : 1.15) and seeds (IZ : 22mm; AI : 1.10) exhibit maximum antifungal activity against *A. niger*.

In other metabolites rich fractions, acetone extracts of both plant parts were found to be more active against *A. niger* (flowers: IZ: 23 mm, AI: 1.15; seeds: IZ: 18mm, AI: 0.72) as compare to other extracts. Earlier, *D. regia* has been reported by a number of workers, with an aim to identify the active principle (s) and their bioefficacy (Thiribhuvanamala and Narasimhan, 1998; Aqil and Ahmad, 2003; Satish et al., 2007, 2009; Dutta et al., 2008).

It may, therefore be concluded that almost all the metabolite rich fractions, flavonoid fractions, anthraquinones fractions and sterol fractions exhibit antifungal activity except aqueous fraction that is rich in sugars. The above findings would prove useful in undertaking further the bioactivity oriented separation of compounds in *D. regia* and can be used as source of indigenous medicine.

REFERENCES

- Ahmad, I., Mahmood, Z. and Mohammad, F. 1998. Screening of some Indian medicinal plants for their antimicrobial properties. *J. Ethnopharmacol.* 62: 183-193.
- Aqil, F. and Ahmad, I. 2003. Broad-spectrum antibacterial and antifungal properties of certain traditionally used Indian medicinal plants. *World J. Microbiol. and Biotech.* 19(6): 653-657.
- Bauer, A. W., Kirby, W. M. M. and Sherris, J. C. 1966. Antibiotic susceptibility testing by a standardized single disk method. *Am. J. Clin. Pathol.* 45: 493-496.
- Bhavanani, S. M. and Ballow, C. H. 2000. New agents for Gram-positive bacteria. *Curr. Opin. Microbiol.* 3: 528-534.
- Clark, A. M. 1996. Natural products as resource for new drugs. *Pharm. Res.* 13: 1133-1141.
- Chung, P. Y., Chung, L. Y., Ngeow, Y. F., Goh, S. F. and Imdiyabing, Z. 2004. Antimicrobial activities of Malaysian plant species. *Pharm. Biol.* 42: 292-300.
- Cordell, G. A. 2000. Biodiversity and drug discovery: a symbiotic relationship. *Phytochemistry.* 55: 463-480.
- Davis, J. 1994. Inactivation of the antibiotics and the dissemination of resistance gene. *Science.* 264: 375-382.
- De Boer, H. J., Kool, A. and Broberg, A. 2005. Antifungal and antibacterial activity of some herbal remedies from Tanzania. *J. Ethnopharmacol.* 96: 461-469.